



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Variations of AIRS CO₂ in the Polar Region

**Xun Jiang¹, Edward Olsen², Jingqian Wang¹,
Thomas Pagano², Luke Chen², and Yuk Yung³**

¹ Department of Earth & Atmospheric Sciences, University of Houston

² Science Division, Jet Propulsion Laboratory, Caltech

³ Division of Geological & Planetary Sciences, Caltech

AIRS Science Team Meeting, April 24-27, 2012



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Overview

- **Motivation**
- **Data**
- **Variability of AIRS CO₂ in the High Latitudes**
Northern Annular Mode, Stratospheric Sudden Warming



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Motivation

- **There are few studies of the CO₂ variability in the polar region due to limited observations.**
- **Variability of CO₂ in the polar region is important for it impacts climate change by modifying the radiation budget and hence the extent of snow and ice.**
- **Global distributed CO₂ retrievals from AIRS can be used to investigate the influence of the large scale dynamics on the polar CO₂.**



National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

AIRS Data Sensitivity in Atmospheric Column

➤ **AIRS Mid-tropospheric CO₂
Sensitivity Peak: 500-300 hPa
(dependent upon latitude)**

Chahine et al. [2005; 2008]






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Influence of Northern Annular Mode on AIRS CO₂



Detrended AIRS CO₂ from Nov to Apr at 60N - 90N (Black)
Detrended & Inverted Arctic Oscillation Index from Nov to Apr (Red)
Correlation Coefficient = 0.74




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sou

Influence of Northern Annular Mode on AIRS CO₂ (Strong Vortex)



AIRS CO₂ averaged in 2005 and 2007 (Nov-Apr)
Positive AO index; Strong Vortex




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared S

Influence of Northern Annular Mode on AIRS CO₂ (Weak Vortex)



AIRS CO₂ averaged in 2006 and 2008 (Nov-Apr)
Negative AO index; Weak Vortex




National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared So

Influence of Northern Annular Mode on AIRS CO₂ (Strong/Weak Vortex Contrast)



AIRS CO₂ Difference
Strong Vortex - Weak Vortex



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder

Influence of Stratospheric Sudden Warming on AIRS CO₂ (SSW of March 2005)

A **stratospheric warming** occurs when the latitudinal **gradient** in 10-hPa zonal-mean temperatures between 85°N-60°N is **positive** for more than 5 days. [WMO]

If the 10-hPa zonal wind at 65°N is concurrently easterly, the warming event is categorized as a “**major warming**”.

Otherwise, the warming event is categorized as “**minor**”.



NCEP 30 hPa mean
temperature during SSW in
Mar 2005



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Influence of Stratospheric Sudden Warming on AIRS CO₂ (EOF Method of Analysis)

Empirical Orthogonal Function (EOF):

The empirical orthogonal function expansion (EOF) is also known as the principal component analysis (PCA), or singular value decomposition method. The essence of EOF is briefly summarized as follows:

$$z(x, t) = \sum_1^n a_k(t) f_k(x)$$

$z(x, t)$: real data

a_k : time series, Principle Component (PC) time series

f_k : modes of spatial pattern, EOFs, Eigenvectors of Covariance Matrix

$$(C = z^T \cdot z / (N-1))$$

Eigenvalue of C: the fraction of variance capture by EOF

EOF1 of GPH captures the Northern Annular Mode (NAM)



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Vertical Structure of the Northern Annular Mode

(Strength of Polar Vortex is Characterized by the NAM index)

- Spatial patterns of NCEP-2 geopotential heights leading mode.

- Vertical structure of the Northern Annular Mode (NAM) index from NCEP-2 geopotential height (EOF1 of GPH)

MARCH 17 - COLLAPSE OF POLAR VORTEX
(coupling of stratosphere and troposphere results)





National Aeronautics and
Space Administration


Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder


Temperature Before/After SSW

NCEP2 Temperature


Before SSW



During SSW



After SSW






National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California


Atmospheric Infrared Sounder

AIRS Mid-tropospheric CO₂ Before and After March 2005 SSW



**Before SSW
of March 2005**

Vortex isolates polar atmosphere
from Mid-lat atmosphere
tropopause lowered



**After SSW
of March 2005**

Polar vortex area shrinks and breaks
Mid-lat higher CO2 concentrations
mix into the polar regions



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Conclusions

- **AIRS CO₂ at high latitudes correlate well with the strength of the northern hemispheric polar vortex in the winter season.**
- **During the Stratospheric Sudden Warming (SSW) event, the polar temperature increases and polar winds switch from westerly to easterly. Polar mid-tropospheric CO₂ concentrations increase after the SSW in March 2005.**



National Aeronautics and
Space Administration

Jet Propulsion Laboratory
California Institute of Technology
Pasadena, California

Atmospheric Infrared Sounder

Thank you!